



TONY KIMMONS, DVM

**4660 Murfreesboro Road
Franklin, TN 37067**

**P.O. Box 369
Arrington, TN 37014**

615-395-4441

615-395-4152: Fax

Adult vaccination chart:

Core Vaccines	Vaccine	Performance & Pleasure horses	Unvaccinated horses or with unknown history	Pregnant mares
	Equine Eastern/Western Encephalomyelitis	Annual, Spring	2 doses 4-6 weeks apart then annually in Spring	Annual, 4-6 weeks prior to foaling
	Tetanus	Annual. Repeat after penetrating wound unless vaccinated within 6 months.	2 doses 4-6 weeks apart then annually	Annual, 4-6 weeks prior to foaling
	Herpesviruses EHV-1 and EHV-4) = Rhinopneumonitis	Every 6 months. In the event you are showing/traveling, it is recommended to give every 3 months, otherwise every 6 months.	Series of 3 at 4-6 weeks apart then every 6 months.	Inactivated EHV-1 vaccine at 3, 5, 7 and 9 months of gestation. EHV-1 and EHV-4 4-6 weeks prior to foaling.
	West Nile Vaccine (Prevenile)	Annual, Spring	One dose, then annually in Spring	Annual, 4-6 weeks prior to foaling.
	Rabies	Annual	One dose then annually	Annual, 4-6 weeks prior to foaling
	Equine Influenza	Intranasal every 6 months	Intranasal every 6 months	Intranasal every 6 months with booster 4-6 weeks prior to foaling.
Optional vaccines	Strangles (intranasal) (Optional)	Optional, depends on risk. Every 6-12 months. WE STRONGLY SUGGEST THIS VACCINATION.	Optional, depends on risk. WE STRONGLY SUGGEST THIS VACCINATION. 2 doses w/ 3 week interval then every 6-12 months.	Optional, depends on risk. WE STRONGLY SUGGEST THIS VACCINATION.
	Potomac Horse Fever (Optional)	Not routine in this area. If traveling to endemic areas, give semi-annually	Not routine in this area. If traveling to endemic areas, give semi-annually	
	Botulism (Optional)	Annual, Spring	3 doses 1 month apart, then annually	Annual, 4-6 weeks prior to foaling